

AD-A127 535

A CLASS OF MULTIVARIATE NEW BETTER THAN USED PROCESSES
(U) ILLINOIS UNIV AT CHICAGO CIRCLE DEPT OF MATHEMATICS
E EL-NEWEINI DEC 82 AFOSR-TR-83-0297 AFOSR-80-0170


1/1

UNCLASSIFIED

F/G 12/1

NL





MICROCOPY RESOLUTION TEST CHART
NATIONAL BUREAU OF STANDARDS-1963-A

(4)

AD A127535

A Class of Multivariate New Better
Than used Processes

by

Emad El-Newehi*

ERIC
S D
APR 20 1983
E

Approved for public release?
Distribution unlimited.

NE

83-04-28-010

A Class of Multivariate New Better
Than used Processes

by

Emad El-Newehi*

December, 1982.

Department of Mathematics, Statistics
and Computer Science
University of Illinois at Chicago
Chicago, Illinois

DTIC
ELECTE
S APR 29 1983 D
E

Research sponsored by the Air Force Office of Scientific Research,
AFSC, USAF, under Grant AFOSR 80-0170.

Key words and phrases: Multivariate new better than used processes
multivariate new better than used distributions, monotone systems,
convolution, upper sets, lower sets.

AMS 1980 subject classifications. Primary 60K10; Secondary 62N05, 62H05.

AIR FORCE OFFICE OF SCIENTIFIC RESEARCH (AFOSR)
NOTICE OF TRANSMISSION TO DTIC
This technical report is approved for release and is
approved for release under the provisions of AFOSR-12.
Distribution is unlimited.
MATTHEW J. KENNEL
Chief, Technical Information Division

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

REPORT DOCUMENTATION PAGE		READ INSTRUCTIONS BEFORE COMPLETING FORM
1. REPORT NUMBER AFOSR-TR- 83-0297	2. GOVT ACCESSION NO. <i>AP-A127535</i>	3. RECIPIENT'S CATALOG NUMBER
4. TITLE (and Subtitle) A CLASS OF MULTIVARIATE NEW BETTER THAN USED PROCESSES		5. TYPE OF REPORT & PERIOD COVERED TECHNICAL
		6. PERFORMING ORG. REPORT NUMBER
7. AUTHOR(s) Emed El-Neweihi		8. CONTRACT OR GRANT NUMBER(s) AFOSR-80-0170
9. PERFORMING ORGANIZATION NAME AND ADDRESS Department of Mathematics, Statistics, and Computer Science, University of Illinois, P.O. Box 4348, Chicago IL 60680		10. PROGRAM ELEMENT, PROJECT, TASK AREA & WORK UNIT NUMBERS PE61102F; 2304/A5
11. CONTROLLING OFFICE NAME AND ADDRESS Mathematical & Information Sciences Directorate Air Force Office of Scientific Research Bolling AFB DC 20332		12. REPORT DATE DEC 82
		13. NUMBER OF PAGES 9
14. MONITORING AGENCY NAME & ADDRESS (if different from Controlling Office)		15. SECURITY CLASS. (of this report) UNCLASSIFIED
		15a. DECLASSIFICATION/DOWNGRADING SCHEDULE
16. DISTRIBUTION STATEMENT (of this Report) Approved for public release; distribution unlimited.		
17. DISTRIBUTION STATEMENT (of the abstract entered in Block 20, if different from Report)		
18. SUPPLEMENTARY NOTES		
19. KEY WORDS (Continue on reverse side if necessary and identify by block number) Multivariate new better than used processes; multivariate new better than used distributions; monotone systems; convolution; upper sets; lower sets.		
20. ABSTRACT (Continue on reverse side if necessary and identify by block number) A multivariate version of the univariate new better than used (NBU) processes, due to El-Neweihi, Proschan and Sethuraman is introduced. Closure properties of this new class of multivariate NBU (MNBU) processes under various reliability operations are obtained. A relationship between the MNBU processes and a well known class of multivariate NBU distributions is described.		

DD FORM 1 JAN 73 1473

UNCLASSIFIED

SECURITY CLASSIFICATION OF THIS PAGE (When Data Entered)

83 04 28 040

Abstract

A multivariate version of the univariate new better than used (NBU) processes, due to El-Newehi, Proschan, and Sethuraman is introduced. Closure properties of this new class of multivariate NBU (MNBU) processes under various reliability operations are obtained. A relationship between the MNBU processes and a well known class of multivariate NBU distributions is described.

Accession For	
NTIS GRA&I	<input checked="checked" type="checkbox"/>
DTIC TAB	<input type="checkbox"/>
Unannounced	<input type="checkbox"/>
Justification	
By	
Distribution/	
Availability Codes	
Dist	Avail and/or Special
A	



1. Introduction.

The theory of multistate coherent systems has motivated several authors to introduce and study special classes of stochastic processes which generalize the well known classes of lifetimes in the two-state reliability theory. In a paper by El-Newehi, Proschan and Sethuraman (1978) a univariate class of new better than used (NBU) processes was introduced. A member in this class can serve as a model to describe the degradation of a multistate component. A multivariate process whose univariate marginals are independent NBU processes can then be used to describe the joint stochastic behavior of a collection of degradable components. However such a model is not adequate when it is unreasonable to assume stochastic independence. The main purpose of this paper is to introduce a more general class of multivariate NBU processes which extends the univariate NBU class due to El-Newehi, Proschan and Sethuraman (1978). The approach used in this paper is similar to the one used by Block and Savits (1981) to introduce a class of multivariate IFRA processes.

A brief summary of the contents of this paper is now given. In section 2 the basic notation, terminology and definitions are given. In section 3 a class of multivariate NBU processes (hereafter referred to as the MNBU class) is defined. Necessary and sufficient conditions for a multivariate process to be a member in the MNBU class are given. Finally in section 4 it is shown that the MNBU class enjoys a number of closure properties which include as special cases some well known preservation properties of the univariate class of NBU lifetimes. Some of these properties are utilized to construct examples of MNBU processes. Using one of the characterizations established in section 3, the MNBU class is shown to be closely related to a well known class of multivariate HOU distributions.

2. Notation, Definitions and Terminology.

The vector $\underline{x} = (x_1, \dots, x_n)$ denotes a point in R^n , the real n -dimensional Euclidean space equipped with its usual metric.

$\underline{y} \leq \underline{x}$ means that $y_i \leq x_i$, $i = 1, \dots, n$

$\underline{y} < \underline{x}$ means that $y_i < x_i$, $i = 1, \dots, n$

A subset $U \subseteq R^n$ is an upper set if $\underline{x} \in U$ and $\underline{x} \leq \underline{y}$ imply that $\underline{y} \in U$.

Given a vector $\underline{x} \in R^n$, $Q_{\underline{x}}$ denotes the set $\{\underline{y}: \underline{x} < \underline{y}\}$.

A function $f: R^n \rightarrow R^m$ is a nondecreasing function if $\underline{x} \leq \underline{y}$ implies that $f(\underline{x}) \leq f(\underline{y})$. A nonincreasing function is similarly defined.

Throughout the remainder of this paper all the random variables (random vectors) considered are assumed to be nonnegative.

3. The MNBU Class: Definition and Characterizations.

First let us recall that a nonnegative random variable T is said to be IBU if

$$P(T > t) \leq P(T > \alpha t)P(T > (1-\alpha)t), \quad (3.1)$$

for every $t \geq 0$ and every $0 \leq \alpha \leq 1$.

In the context of reliability theory the random variable T usually represents the lifetime (functioning time) of a component that can be in either of two states "functioning" (denoted by 1) and "failed" (denoted by 0). In a paper by El-Newehi, Proschan and Sethuraman (1970) a class of univariate IBU stochastic process was defined as follows: A nonnegative, nonincreasing and right-continuous stochastic process $\{X(t): t \geq 0\}$

whose state space is $\{0,1,\dots,M\}$ is said to be NBU if

$$T_j = \inf \{t: X(t) \notin \{j+1,\dots,M\}\} \quad (3.2)$$

is NBU random variable for every $j \in \{0,1,\dots,M\}$. Such a process was introduced to describe the stochastic behavior of a multistate component which starts at time 0 in state M (perfect functioning) and deteriorates, as time passes, to lower states until finally state 0 (complete failure) is reached. The definition of the NBU process can be easily extended to the case in which the state space S is any subset of $[0,\infty)$ by simply requiring that the random variable

$$T_a = \inf \{t: X(t) \notin (a,\infty)\} \quad (3.3)$$

is NBU for every $a \geq 0$.

A natural multivariate extension of (3.3) is now used to define the MNBU class of stochastic processes.

Definition 3.1. A vector-valued stochastic process $\{\underline{X}(t) = (X_1(t), \dots, X_n(t)) : t \geq 0\}$ is said to be a MNBU process if it is nonnegative, nonincreasing, right-continuous and the random variable

$$T_U = \inf \{t: \underline{X}(t) \notin U\} \quad (3.4)$$

is NBU for every open upper set $U \subseteq \mathbb{R}^n$.

The class of all MNBU processes (of all dimensions) is called the MNBU class. Clearly the MNBU class includes, when $n=1$, the univariate NBU class due to El-Newehi, Proschan and Sethuraman (1978). Also in a recent paper by Block and Savits (1981), the authors define a multivariate class of IFRA processes by requiring the random variable in (3.4) to be

IFRA. Clearly their class is a subclass of the MNBU class.

Remark 3.2. In view of (3.1) and the right-continuity of the process $\{\underline{X}(t): t \geq 0\}$, condition (3.4) is equivalent to

$$P(\underline{X}(t) \in U) \leq P(\underline{X}(\alpha t) \in U)P(\underline{X}((1-\alpha)t) \in U), \quad (3.5)$$

for every $t \geq 0$, $0 \leq \alpha \leq 1$ and every open upper set $U \subseteq \mathbb{R}^n$.

In the remainder of this section two characterizations of the MNBU class are given. These characterizations are used in section 4 to establish properties of the MNBU class and relate it to a well known class of multivariate NBU distributions. First we need the following lemma whose proof is essentially due to Esary, Proschan and Walkup (1967)(see also Marshall and Shaked (1982)).

Lemma 3.3. Let $\{\underline{X}(t): t \geq 0\}$ be a stochastic process. Then condition (3.5) is equivalent to

$$P(\underline{X}(t) \in U) \leq p(\underline{X}(\alpha t) \in U)P(\underline{X}((1-\alpha)t) \in U) \quad (3.6)$$

for every $t \geq 0$, $0 \leq \alpha \leq 1$ and every upper Borel set $U \subseteq \mathbb{R}^n$.

The following theorem gives a necessary and sufficient condition for a nonnegative, nonincreasing and right-continuous stochastic process $\{\underline{X}(t): t \geq 0\}$ to be a MNBU process.

Theorem 3.4. The process $\{\underline{X}(t): t \geq 0\}$ is a MNBU process if and only if

$$E[\phi(\underline{X}(t))] \leq E[\phi(\underline{X}(\alpha t))] E[\phi^{1-\alpha}(\underline{X}((1-\alpha)t))], \quad (3.7)$$

for every nonnegative, nondecreasing and Borel measurable function $\phi: \mathbb{R}^n \rightarrow \mathbb{R}$ and every $t \geq 0$, $0 \leq \alpha \leq 1$.

Proof. Obviously (3.7) implies (3.5) by taking $\phi = I_U$, where $U \subseteq R^n$ is an open upper set. Now assume $\{X(t): t \geq 0\}$ is a MINBU process. In view of (3.6), condition (3.7) is satisfied for all $\phi = aI_U$, where $a \geq 0$ and $U \subseteq R^n$ is an upper Borel set. Let $\phi = \sum_{i=1}^m \phi_i$, where $\phi_i = a_i I_{U_i}$, $a_i \geq 0$, $U_i \subseteq R^n$ is an upper Borel set, $i = 1, \dots, m$ and $m \geq 1$. By Holder inequality for sequences and Minkowski inequality for the L_α -norm, $0 < \alpha < 1$, we get

$$\begin{aligned} E[\phi(X(t))] &\leq \sum_{i=1}^m E[\phi_i^\alpha(X(\alpha t))] E[\phi_i^{1-\alpha}(X((1-\alpha)t))] \\ &\leq \left(\sum_{i=1}^m [E[\phi_i^\alpha(X(\alpha t))]]^{1/\alpha} \right) \left(\sum_{i=1}^m [E[\phi_i^{1-\alpha}(X((1-\alpha)t))]]^{1/(1-\alpha)} \right)^{1-\alpha} \\ &\leq E[\phi^\alpha(X(\alpha t))] E[\phi^{1-\alpha}(X((1-\alpha)t))]. \end{aligned}$$

The general result now follows by taking monotone sequences of such functions and passing to the limit.

Remark 3.5. In a recent paper by Marshall and Shaked (1982) the authors established a similar characterization for their class of multivariate NBU distributions. However they used a different method to extend the validity of the inequality in (3.7) from indicator functions to nonnegative linear combinations of indicator functions.

The second characterization for the MINBU class, which is useful in relating it to a multivariate concept of NBU distributions, follows from the following lemma which is due to Block and Savits (1980). We provide here a simple and new proof.

Lemma 3.6. Let $U \subseteq R^n$ be an open upper set. Then $U = \bigcup_{i=1}^{\infty} Q_{\underline{x}^i}$, where $\underline{x}^i \in R^n$, $i = 1, 2, \dots$.

Proof. Let \underline{x} be an arbitrary element in U . Since U is open there exists a $\underline{y} \in U$ such that $\underline{y} < \underline{x}$ and y_i is rational, $i = 1, 2, \dots, n$. The result now follows by observing that $\underline{x} \in Q_{\underline{y}} \subseteq U$ and that the set of points in R^n with rational coordinates is countable.

Theorem 3.7. The following condition is equivalent to (3.5),

$$P(\underline{X}(t) \in U) \leq P(\underline{X}(\alpha t) \in U)P(\underline{X}((1-\alpha)t) \in U),$$

for all $t \geq 0$, $0 \leq \alpha \leq 1$ and all U of the form $\bigcup_{i=1}^p Q_{\underline{x}^i}$ where $\underline{x}^i \in R^n$, $i = 1, \dots, p$, p is a positive integer.

Proof. Obviously (3.5) implies (3.8). Now assume (3.8) is true and let U be an open upper set. By lemma 3.6 there exists a sequence $U_1 \subseteq U_2 \subseteq \dots$ increasing to U and (3.8) is true for each U_i . The result now follows by passing to the limit.

4. Closure Properties, Examples and a Related Class of Multivariate NBU Distributions.

Closure properties of the MINBU class, which correspond to common reliability operations, are established in this section. Such properties are then utilized to construct and identify members of the MINBU class.

In the following theorem three fundamental closure properties of the MINBU class are established.

Theorem 4.1. The following properties hold for the MINBU class:

(P) Let $\underline{X}(t) = (X_1(t), \dots, X_n(t))$; $t \geq 0$ be a MINBU process and let $f: R^n \rightarrow R^n$ be a nonnegative, nondecreasing and left-continuous function. Then $\{f(\underline{X}(t)); t \geq 0\}$ is a MINBU process.

(P2) Let $\{\underline{X}(t) = (X_1(t), \dots, X_n(t)): t \geq 0\}$ and $\{Y(t) = (Y_1(t), \dots, Y_m(t)): t \geq 0\}$ be two independent MIBU processes. Then $\{(\underline{X}(t); Y(t)): t \geq 0\}$ is a $(m+n)$ -dimensional MIBU process.

(P3) Let $\{\underline{X}_n(t): t \geq 0\}$, $n=1,2,\dots$ be a sequence of k -dimensional MIBU processes and $\underline{X}_n(t) \leq \underline{X}(t)$ for every t , where $\{\underline{X}(t): t \geq 0\}$ is nonnegative nonincreasing and right-continuous process. Then $\{\underline{X}(t): t \geq 0\}$ is a MIBU process.

Proof. (P1) The proof is obvious in view of theorem (3.4).

(P2) The proof is similar to the one given by Marshall and Shaked (1982), pages 202-203, to establish a similar property for their class of multivariate HBU distributions. The details are therefore omitted.

(P3) Let $U \subseteq R^k$ be an open upper set. Let $\phi_m: R^k \rightarrow R$ be a sequence of continuous, bounded and nondecreasing functions such that $\phi_m \rightarrow I_U$ pointwise. By theorem 3.4 we have

$$E[\phi_m(\underline{X}(t))] \leq E[\phi_m^\alpha(\underline{X}_n(\alpha t))] \leq E[\phi_m^{1-\alpha}(\underline{X}_n((1-\alpha)t))],$$

The result now follows by taking limits as $n \rightarrow \infty$ then as $m \rightarrow \infty$.

Remark 4.2. The assumption of left-continuity in (P1) is not needed when the state space for the process $\{\underline{X}(t): t \geq 0\}$ is finite.

The following properties for the MIBU class are immediate consequences of the fundamental properties (P1) and (P2).

Corollary 4.3. The following properties hold for the MIBU class:

(P3) Let $\{X_i(t): t \geq 0\}$, $i=1,\dots,n$ be independent univariate HBU processes. Then $\{\underline{X}(t) = (X_1(t), \dots, X_n(t)): t \geq 0\}$ is a MIBU process.

(P4) Let $\{\underline{X}(t): t \geq 0\}$ be a n -dimensional MIBU process. Then

$\{X(t) = (X_{i_1}(t), \dots, X_{i_k}(t)): t \geq 0\}$ is a k -dimensional MIBU process, where $1 \leq k \leq n$ and $1 \leq i_1 < \dots < i_k \leq n$.

(P5) Let $\{\underline{X}(t) = (X_1(t), \dots, X_n(t)) : t \geq 0\}$ be a MIBU process and let $\phi_i: \mathbb{R}^n \rightarrow \mathbb{R}$ be nonnegative, nondecreasing and left-continuous function, $i=1, \dots, m$. Then $\{(\phi_1(\underline{X}(t)), \dots, \phi_m(\underline{X}(t))) : t \geq 0\}$ is a MIBU process.

(P6) Let $\{\underline{X}_i(t) : t \geq 0\}$, $i=1, \dots, k$ be k independent MIBU processes of the same dimension. Then $\{\sum_{i=1}^k \underline{X}_i(t) : t \geq 0\}$ is a MIBU process.

Proof. The proof is obvious and is therefore omitted.

Remark 4.4. It should be noted that the generalized MIBU closure theorem due to El-Howeishi, Proschan and Sethuraman (1973) is a special case of (P5). Also observe that (P6) asserts the closure of the MIBU class under convolution.

The following theorem relates the MIBU class to a well known class of multivariate MIBU random vectors, namely the Class $C = \{\underline{I} = (T_1, \dots, T_n) : \underline{I} \text{ is nonnegative and } \tau(\underline{I}) \text{ is MIBU for every monotone life function } \tau, n \text{ is arbitrary positive integer. (recall that } \tau(\underline{I}) \text{ has the form } \max_{1 \leq j \leq p} \min_{i \in A_j} T_i, \text{ where } \bigcup_{j=1}^p A_j \subseteq \{1, \dots, n\})\}$.

Theorem 4.5. The process $\{\underline{X}(t) : t \geq 0\}$ is a MIBU process if and only if every finite collection of $\{T_{ix} : 1 \leq i \leq n, x \in \mathbb{R}\}$ is in C , where $T_{ix} = \inf\{t : X_i(t) \leq x\}$.

Proof. The proof follows readily by theorem 3.7 and the simple observation that for any two upper Borel sets U_1, U_2 , $T_{U_1 \cup U_2} = \max(T_{U_1}, T_{U_2})$ and $T_{U_1 \cap U_2} = \min(T_{U_1}, T_{U_2})$. The simple details of the proof are left to the reader.

Remark 4.6. The above theorem is essentially the analogue of theorem 2.4 of Block and Savits (1981).

The above properties of the MIBU class and its relationship to the class C of multivariate IDU random vectors can be utilized to construct many examples of MIBU processes. For instance if \underline{I} belongs to C then $\{(I_{(T_1 > t)}, \dots, I_{(T_n > t)}): t \geq 0\}$ is a MIBU process. Also multivariate processes whose coordinates are monotone functions of independent univariate IDU processes are MIBU processes. The following example motivated by the theory of multistate coherent systems illustrates the construction and identification of members of the MIBU class.

Example 4.7. Consider a system of n binary independent components. Assume we have $M-1$ spares for each of the n components. A failed component is instantaneously replaced by one of its spares. All the lifetimes are assumed to be independent IDU random variables. When the original component i is functioning (and none of the spares has been used), we consider that component i is in state 1. Upon failure of the original component one of its spares is used and so the component now enters state $M-1$, etc. Let $\{\underline{X}(t) = (X_1(t), \dots, X_n(t)): t \geq 0\}$ be the process describing the states of the n components as time passes. Then $\{\underline{X}(t): t \geq 0\}$ is a MIBU process. Moreover if the n components are forming a coherent system whose cut sets are C_1, \dots, C_r and $Y_j(t) = \max_{i \in C_j} (X_i(t))$, $j=1, \dots, r$, then $\{(Y_1(t), \dots, Y_r(t)): t \geq 0\}$ is also a MIBU process.

REFERENCES

1. Block, H.W. and Savits, T.H. (1980). Multivariate increasing failure rate average distributions. Ann. Probability, 8, 793-801.
2. Block, H.W. and Savits, T.H. (1981). Multidimensional IFRA processes. Ann. Probability, 9, 162-166.
3. El-Newehi, E., Proschan, F. and Sethuraman, J. (1978). Multistate coherent systems. J. Appl. Prob., 15, 575 - 588.
4. Esary, J.D., Proschan, F. and Walkup, D.W. (1967). Association of random variables with applications. Ann. Math. Statist., 38, 1466-1474.
5. Marshall, A.W. and Shaked, M. (1982). A class of multivariate new better than used distributions. Ann. Probability, 10, 259-264.

END

FILMED

6-83

DTIC